

I. J. SCHOENBERG

The Landau problem I.

The case of motions on sets

Dedicated to N. G. de Bruijn on the occasion of his 60th birthday

Communicated by Prof J. H. van Lint at the meeting of December 17, 1977

*Department of Mathematics, United States Military Academy, West Point,
New York 10996*

1. STATEMENT OF THE PROBLEM

Let $f(t)$ be real-valued, $-\infty < t < \infty$. As we think of t as time, we denote its derivatives by $\dot{f}(t)$ and $\ddot{f}(t)$. We assume that $f \in C^1(\mathbb{R})$, hence $\dot{f} \in C(\mathbb{R})$, and that $\dot{f}(t)$ is piecewise continuous with discontinuities only of the first kind. Using the supremum norm on \mathbb{R} , we have the well known inequality of Landau

$$(1) \quad \|\dot{f}\| \leq \sqrt{2 \cdot \|f\| \cdot \|\ddot{f}\|}, \text{ (See [1], [2]).}$$

If $a > 0$ and $f(t)$ satisfies the inequalities

$$(2) \quad -a \leq f(t) \leq a \text{ for all } t,$$

then (1) shows that

$$(3) \quad \|\dot{f}\| \leq \sqrt{2a} \sqrt{\|\ddot{f}\|}.$$

Here $\sqrt{2a}$ is the best constant, as shown by the special motion $\dot{f}(t) = 4a(t - t^2)$ if $0 \leq t \leq 1$, which is extended to all real t by the functional equation $\dot{f}(t+1) = -\dot{f}(t)$, for all real t , with the result that $\|\dot{f}\| = a$, $\|\ddot{f}\| = 4a$, $\|\ddot{\dot{f}}\| = 8a$. This motion produces the equality sign in (3).

We may interpret (2) by saying that $f(t)$ describes a motion in the interval

$$(4) \quad I_a = \{-a \leq x \leq a\}.$$

In terms of the functional

$$(5) \quad F(f) = \frac{\|f\|}{\sqrt{\|f\|^2}}$$

we may restate the result (3) by writing that

$$(6) \quad \sup_f F(f) = \sqrt{2a}$$

for all motions f within the interval I_a .

It seems rather natural to replace the interval (4) by a given *closed and connected set* S of the complex plane \mathbb{C} , or belonging to a finite-dimensional euclidean space. Accordingly, we consider the class of motions

$$(7) \quad (S) = \{f(t); f(t) \in S \text{ for all } t, f(t) \neq \text{constant}\}.$$

Notice that we have excluded the trivial case when $f(t) = \text{constant}$ for all t , when $F(f) = 0/0$ is undefined.

We can now formulate

LANDAU'S PROBLEM FOR THE SET S . *To determine the quantity*

$$(8) \quad L(S) = \sup_{f \in (S)} F(f).$$

We call $L(S)$ the Landau constant of the set S .

2. A FEW SPECIAL SETS S WHEN $L(S)$ IS KNOWN

In terms of (1.4) and the definition (1.8), we may restate the result (1.6) by writing that

$$(1) \quad L(I_a) = \sqrt{2a}.$$

Here are a few trivial further examples. If $S = \{0 \leq x < \infty\}$, then $f(t) = t^2 \in (S)$, and $\|f\| = \infty$, $\|f'\| = 2$. Therefore $L(S) = \infty/2 = \infty$. If $S = \mathbb{R}$, then $f(t) = t \in (S)$, and $L(S) = 1/0 = \infty$.

An obvious and fundamental property of the Landau constant is the following

MONOTONICITY PROPERTY. *If S_1 and S_2 are two sets, then*

$$(2) \quad S_1 \subset S_2 \text{ implies that } L(S_1) \leq L(S_2).$$

A few further non-trivial examples when $L(S)$ is known are the following. If S is the *circular disc*

$$(3) \quad D_a = \{|z| \leq a\},$$

then $I_a \subset D_a$ implies, by (2), that $L(I_a) \leq L(D_a)$. As a matter of fact we have here equality, so that

$$(4) \quad L(D_a) = \sqrt{2a}.$$

For the *circle*

$$(5) \quad C_a = \{|z| = a\}$$

we will show that

$$(6) \quad L(C_a) = \sqrt{a}.$$

It seems worthwhile to look at the two results (1) and (6) and compare their respective extremizing motions. Let $\mathcal{E}_2(t)$ be the so-called *quadratic Euler spline* (See [2, Section 2]) defined by

$$(7) \quad \mathcal{E}_2(t) = 1 - 4t^2 \text{ if } -\frac{1}{2} \leq t \leq \frac{1}{2}, \text{ and } \mathcal{E}_2(t+1) = -\mathcal{E}_2(t) \text{ for all } t.$$

We found above that $\tilde{f}(t) = a\mathcal{E}_2(t) \in (I_a)$ and $F(\tilde{f}) = \sqrt{2a}$. For $S = C_a$ the extremizing motion is the *uniform rotation*

$$\tilde{f}(t) = ae^{it}.$$

Indeed, here $\|\dot{\tilde{f}}\| = a$, $\|\ddot{\tilde{f}}\| = a$, hence $F(\tilde{f}) = \sqrt{a}$, and this is the extremizing motion in view of (6).

Let now S be the *circular ring*

$$(8) \quad R_{a,b} = \{b \leq |z| \leq a\}, \quad (0 < b < a).$$

Here we shall find that *

$$(9) \quad L(R_{a,b}) = \sqrt{a + \sqrt{a^2 - b^2}}.$$

We would expect to obtain from this the result (6) when $b = a$, and (4) when $b = 0$.

Our last special example is the *solid shell*

$$(10) \quad SoSh_{a,b} = \{b \leq \sqrt{x^2 + y^2 + z^2} \leq a\}, \quad (0 < b < a).$$

As this will yield most of the previous results, we formulate it as

THEOREM 1. *For the solid shell (10) we have that*

$$(11) \quad L(SoSh_{a,b}) = \sqrt{a + \sqrt{a^2 - b^2}}.$$

Actually Theorem 1, with the same value for the Landau constant, holds for the solid spherical shell in n dimensions, $n \geq 2$. Observe again, that $R_{a,b} \subset SoSh_{a,b}$, while their Landau constants are equal by (9) and (11).

Of an entirely different nature is

THEOREM 2. *If the set S is closed, bounded and convex, then*

$$(12) \quad L(S) = \sqrt{\text{diameter of } S}.$$

* This was first shown in [3]. In Section 5 below we first derive the stronger relation (11) which is somewhat easier to prove than (9). Theorem 2 below was stated without proof in [3].

Clearly Theorem 2 implies as special cases the results (1) and (4). Again Theorem 2, and its proof below, remains valid, as it stands, in any finite-dimensional euclidean space.

3. A LEMMA

The lemma to be now discussed, will show that an extremizing function $\tilde{f}(t)$, i.e. producing the equality $F(\tilde{f}) = L(S)$, may also be characterized as a solution of a problem of Optimal Control.

LEMMA 1. *If the motion*

$$(1) \quad \tilde{f}(t) \in (S)$$

has the

PROPERTY A. *If*

$$(2) \quad f(t) \in (S) \text{ and } \|\dot{f}\| \leq \|\ddot{f}\|$$

then

$$(3) \quad \|\dot{f}\| \leq \|\ddot{f}\|,$$

then \tilde{f} also has the

PROPERTY B.

$$(4) \quad F(\tilde{f}) = L(S),$$

and conversely, if \tilde{f} enjoys the Property B, then \tilde{f} also has the Property A.

PROOF: 1. *Property A implies Property B.* We assume (1) and Property A and we are to establish (4). If $f \in (S)$, then for the motion $g(t) = f(t) \sqrt{\|\ddot{f}\| / \|\dot{f}\|}$ we find that $g(t) \in (S)$ and also that

$$\|\dot{g}\| = \|\dot{f}\| (\|\ddot{f}\| / \|\dot{f}\|), \text{ hence } \|\dot{g}\| = \|\ddot{f}\|.$$

Hence $g(t)$ satisfies the assumptions (2), and by Property A we conclude that $\|\dot{g}\| \leq \|\ddot{f}\|$, and by the definition of $g(t)$, that

$$\|\dot{f}\| \sqrt{\|\ddot{f}\| / \|\dot{f}\|} \leq \|\ddot{f}\|, \text{ and so } F(f) \leq F(\tilde{f}).$$

But then

$$L(S) = \sup_{f \in (S)} F(f) \leq F(\tilde{f}).$$

From $F(\tilde{f}) \leq L(S)$ we now conclude that (4) holds.

2. *Property B implies Property A.* Now we assume (1) and (4), and that f satisfies (2), and we are to show that (3) holds. From (4) we obtain

that $F(f) \leq F(\dot{f})$, or $\|\dot{f}\|/\sqrt{\|\dot{f}\|} \leq \|\ddot{f}\|/\sqrt{\|\ddot{f}\|}$, whence

$$\|\dot{f}\| \leq \|\ddot{f}\| \sqrt{\|\dot{f}\|/\|\ddot{f}\|} \leq \|\ddot{f}\|$$

in view of the second inequality (2), and (3) is established. Lemma 1 was stated without proof in [3].

4. A PROOF OF THEOREM 2

Let A and B be points of S such that

$$(1) \quad |B - A| = d = \text{diameter of } S, \quad (A \in S, B \in S).$$

From the relations (2.7) we find that $-1 \leq \mathcal{E}_2(t) \leq 1$ for all t , and that $\mathcal{E}_2(t)$ represents a to-and-fro motion on $[-1, 1]$. We now define on the segment $[A, B]$ the motion

$$(2) \quad \begin{aligned} \tilde{f}(t) &= \frac{1}{2}A(1 - \mathcal{E}_2(t)) + \frac{1}{2}B(1 + \mathcal{E}_2(t)) \\ &= \frac{1}{2}(B + A) + \frac{1}{2}(B - A)\mathcal{E}_2(t). \end{aligned}$$

From $\|\mathcal{E}_2\| = 4$, $\|\ddot{\mathcal{E}}_2\| = 8$, and (1), we find that

$$(3) \quad \|\dot{\tilde{f}}\| = 2d, \quad \|\ddot{\tilde{f}}\| = 4d,$$

and therefore

$$(4) \quad F(\tilde{f}) = \sqrt{d}.$$

We wish to prove that $F(\tilde{f}) = L(S)$, which is the relation (4) of Property B of Lemma 1. By Lemma 1 it suffices to prove the following:

If $f \in (S)$ such that

$$(5) \quad \|\dot{f}\| \leq \|\ddot{f}\| = 4d,$$

then we also have

$$(6) \quad \|\dot{f}\| \leq \|\ddot{f}\| = 2d.$$

This we do as follows: We consider an arbitrary but fixed t_0 such that $\dot{f}(t_0) \neq 0$, and let L be a fixed straight line such that

$$(7) \quad L \text{ is parallel to the vector } \dot{f}(t_0).^*$$

Let the segment $[P, Q] \subset L$ be the orthogonal projection of S onto the line L . Clearly

$$(8) \quad |Q - P| \leq |A - B| = d.$$

Let $f_p(t)$ denote the orthogonal projection of the motion $f(t)$ onto L . By (5) we have

$$\|\dot{f}_p\| \leq \|\dot{f}\| \leq \|\ddot{f}\| = 4d$$

* The reader is asked to sketch a diagram.

and by (8) and Landau's inequality (1.1) we find that

$$\|\dot{f}_p\| \leq \sqrt{|Q-P|} \cdot \sqrt{\|\dot{f}_p\|} \leq \sqrt{d} \cdot \sqrt{4d} = 2d.$$

Now (7) implies that

$$|\dot{f}(t_0)| = |\dot{f}_p(t_0)| \leq \|\dot{f}_p\| \leq 2d.$$

and therefore $\|\dot{f}\| \leq 2d$, since t_0 was arbitrary. This proves (6) and our proof is complete.

REMARK. Our proof shows that for Theorem 2 to be valid, the set S need not be convex. All that is needed is that (1) should hold, and that $[A, B] \subset S$. These assumptions already imply that $L(S) = L([A, B])$. It does seem curious that we can not use the greater freedom of motions in S to increase the value \sqrt{d} of the Landau constant.

5. A PROOF OF THEOREM 1

We now turn to the solid shell

$$(1) \quad S = S_{\text{osh}} = \{b \leq \sqrt{x^2 + y^2 + z^2} \leq a\}, \quad (0 < b < a).$$

Just as in the case of the convex set A we identified the motion (4.2) as the motion that maximizes $F(f)$ among all motions within S , so we will now construct a motion again to be denoted by $\tilde{f}(t)$ ($t \in \mathbb{R}$), that will maximize $F(f)$ for the solid shell (1).




Fig.1

Fig. 1 shows the ring $R_{a,b}$ obtained by intersecting the solid shell with the plane $z=0$. We consider the parabolic arc $\Pi = \widehat{ABA'}$ having its vertex B on the circle C_a , and tangent to C_b at the points A, A' . It will aid our computations if we first show how Π can be constructed. Erect EF perpendicular to OB and intersecting C_a at F . Mark the two points D and V such that

$$(2) \quad DB = BV = EF,$$

and erect the perpendicular to OB at D , intersecting C_b at A and A' . Join V to A and to A' . We claim that

$$(3) \quad \angle OAV = \angle OA'V = 90^\circ.$$

PROOF: From the triangle OEF we find that $(EF)^2 = a^2 - b^2$. However, by (2)

$$OD = a - DB = a - EF, \quad OV = a + BV = a + EF,$$

and so $OD \cdot OV = a^2 - (EF)^2 = a^2 - (a^2 - b^2) = b^2 = (OA')^2$, or $OD \cdot OV = (OA')^2$. This is a characteristic relation for a right-angled triangle and establishes (3).

Incidentally, we have shown that in fig. 1 we have

$$(4) \quad OV = a + \sqrt{a^2 - b^2}.$$

From $DB = BV$ it also follows that the parabolic arc Π with vertex at B and passing through A and A' , is also tangent to C_b at A and A' .

Next we consider on the arc $\Pi = \widehat{ABA'}$ a motion $\tilde{f}(t)$ ($\alpha \leq t \leq \beta$), which we call *Galilean*; by this we mean that the point $\tilde{f}(t)$ describes the arc Π , $\tilde{f}(\alpha) = A$, $\tilde{f}(\beta) = A'$, and that the point $\tilde{f}(t)$ has a *constant acceleration* $\ddot{\tilde{f}}$ which must, of course, be real and negative. For this motion the modulus $|\dot{\tilde{f}}(t)|$ of the velocity vector assumes its maximal value at the endpoints A and A' . Let $v = |\dot{\tilde{f}}(\alpha)| = |\dot{\tilde{f}}(\beta)|$ denote this maximal value. Along Π we therefore have

$$(5) \quad F(\tilde{f}) = v / \sqrt{|\ddot{\tilde{f}}|}.$$

This functional being invariant if we change the scale in t , without loss of generality we may assume $\alpha = 0$, $\beta = 1$, and now

$$(6) \quad \tilde{f}(t), \quad (0 \leq t \leq 1)$$

represents the Galilean motion along the arc Π .

The quantity $F(\tilde{f})$ is of dimension $LT^{-1}(LT^{-2})^{-1/2} = L^{1/2}$. Let us show that *in terms of fig. 1 we have*

$$(7) \quad F(\tilde{f}) = \sqrt{OV}.$$

PROOF: It is well known that a material particle projected vertically upward with the initial velocity v_0 , will reach a maximal height h given by the relation

$$(8) \quad v_0^2 = 2gh.$$

Using (5), the relation (7), to be established, amounts to

$$(9) \quad v^2 = |\ddot{f}| \cdot OV.$$

The motion $\dot{f}(t)$ may be regarded as a motion in the field of constant acceleration $\ddot{f} = -|\ddot{f}|$, while its initial "vertical" velocity is $v_0 = v \sin \gamma$, where $\gamma = \angle DAV$. By (8) we have

$$(v \sin \gamma)^2 = 2|\ddot{f}| \cdot DB = \ddot{f} \cdot DV,$$

in view of (2). It follows that

$$v^2 = |\ddot{f}| \frac{DV}{(\sin \gamma)^2} = |\ddot{f}| \cdot \frac{AV}{\sin \gamma} = |\ddot{f}| \cdot OV$$

proving (9).

We now extend the motion (6) to all real t by the functional equation

$$(10) \quad \dot{f}(t+1) = e^{2i\gamma} \dot{f}(t) \text{ for all real } t.$$

This amounts to successive rigid rotations of the motion on Π through the angle $2\gamma = \angle AOA'$, and shows that the curve Γ described by the extended motion $\dot{f}(t)$ ($t \in \mathbb{R}$), is an infinite sequence of parabolic arcs

$$\dots "II'IIII'I'II" \dots,$$

all congruent to II . Therefore (10) defines a motion $\dot{f}(t)$ having the properties

$$(11) \quad \dot{f}(t) \in (R_{a,b}),$$

$$(12) \quad \dot{f}(t) \in C^1(\mathbb{R}),$$

$$(13) \quad F(\dot{f}) = \sqrt{OV} = \sqrt{a + \sqrt{a^2 - b^2}},$$

the last by (7) and (4).

Notice that the acceleration vector $\ddot{f}(t)$ is piecewise constant, with discontinuities for all integer values of t , while for all other values of t , its modulus is constant, in fact

$$(14) \quad |\ddot{f}(t)| = |\ddot{f}|.$$

After these extensive preliminaries, we can pass to a proof of Theorem 1. Let $f(t)$ be a motion within the solid shell (1) subject to the condition

$$(15) \quad \|\dot{f}\| \leq \|\ddot{f}\| = -\ddot{f},$$

and we are to show that this implies that

$$(16) \quad \|\dot{f}\| \leq \|\dot{\tilde{f}}\| = |\dot{\tilde{f}}(0)| = v.$$

By lemma 1 this would imply that $L(S) = F(\tilde{f})$, and prove Theorem 1 that

$$(17) \quad L(Sh_{a,b}) = F(\tilde{f}) = \sqrt{a + \sqrt{a^2 - b^2}}.$$

To prove (16) we assume that for some t_0 we have $|\dot{\tilde{f}}(t_0)| > v$ and get a contradiction. We may as well assume that $t_0 = 0$, hence

$$(18) \quad |\dot{\tilde{f}}(0)| > v = |\dot{f}(0)|.$$

We begin by rotating the motion $f(t)$ rigidly about 0, hence within the shell S , so that the vector

$$(19) \quad \dot{f}(0) \text{ is parallel to the vector } \dot{\tilde{f}}(0).$$

If $\dot{f}(0)$ is in the half-space containing the arc Π and bounded by the vertical plane π that intersects the horizontal plane xOy in the line A_1OA_3 , we do nothing further. However, if $\dot{f}(0)$ is not in the half-space, then we reflect the entire motion $f(t)$ with respect to π , so as to have now $\dot{f}(0)$ in that half-space; notice that (19) still holds after this reflection. Let $OG \perp A_1A_3$, hence $OG \perp \pi$. Now we rotate the motion $f(t)$ rigidly about the axis of rotation OG so that $\dot{f}(0)$ falls with the quadrant A_1OG . This places $\dot{f}(0)$ within the (partly) curvilinear quadrilateral

$$(20) \quad Q = AA_1H_1H.$$

Assuming that

$$(21) \quad 2\gamma \geq \frac{\pi}{2},$$

which is the case of fig. 1, we find that

$$(22) \quad \dot{f}(O) \in Q' = AA_1A_2A',$$

and it follows that

$$(23) \quad \text{Re } \dot{f}(O) \geq OD.$$

Notice that (19) still holds. We may even assume that the vector

$$(24) \quad \dot{\tilde{f}}(O) \text{ is parallel and points in the same direction as } \dot{\tilde{f}}(O).$$

For if not we replace $f(t)$ by $f(-t)$, thereby reversing the direction of $\dot{f}(O)$.

We now consider the motions $\tilde{f}_p(t)$ and $f_p(t)$ obtained by projecting orthogonally on Ox the motions $\tilde{f}(t)$ and $f(t)$, respectively. Clearly

$$(25) \quad \tilde{f}_p(t) = \text{Re } \tilde{f}(t), \quad f_p(t) = \text{Re } f(t).$$

Let us now compare these rectilinear motions during the time interval from $t=0$ to $t=\frac{1}{2}$. The motion $\tilde{f}_p(t)$ starts from D , at $t=0$, with initial velocity $\dot{\tilde{f}}_p(O)=v \sin \gamma$, and, while uniformly decelerated with constant deceleration $|\ddot{\tilde{f}}|$, just manages to arrive at B , at $t=\frac{1}{2}$, with zero velocity. The motion $f_p(t)$ starts at $\text{Re } f(O)$, ahead of $\tilde{f}_p(O)$, because of (23), with a larger initial velocity, because of (24) and (18), and moves during $0 \leq t \leq \frac{1}{2}$ with an acceleration $\ddot{f}_p(t)$ that nowhere exceeds in modulus the value $|\ddot{\tilde{f}}|$, because of (15). It follows that the point $f_p(t)$ must have reached the point B , for some $t < \frac{1}{2}$, with a positive velocity. This being cinematically impossible, we have obtained the desired contradiction.

There is a slight additional difficulty if (21) does not hold, but rather

$$(26) \quad 2\gamma < \frac{\pi}{2}.$$

Indeed now we would have in fig. 1 that

$$Q = AA_1H_1H \supset Q' = AA_1A_2A'.$$

It follows that while $f(O)$ is in Q , $f(O)$ need not be in Q' . In this case, let δ denote the positive angle $\delta = \angle A'O f(O)$. For the last time we rotate the motion $f(t)$ rigidly clockwise about Oz by the angle δ . As a result the new position of $f(O)$ lies on the segment $[A', A_2]$, and our previous arguments will apply. Indeed observe that (23) will now hold, and that the value of $\dot{f}_p(O)$ has only increased by the last rotation.

REMARK. In fig. 1 we let $b \rightarrow 0+$. As a result we find that

$$\lim_{b \rightarrow 0+} \tilde{f}(t) = a \cdot \mathcal{E}_2(t),$$

uniformly in every finite interval of the t -axis: The extremizing motion for the solid shell converges to the extremizing motion of $I_a = [-a, a]$.

6. COROLLARIES OF THEOREMS 1 AND 2

It was already mentioned that the relations (2.1) and (2.4) are special examples of Theorem 2.

PROOF OF (2.9). From $R_{a,b} \subset \text{SoSh}_{a,b}$, the monotonicity property (2.2) and Theorem 1, we obtain that

$$L(R_{a,b}) \leq \sqrt{a + \sqrt{a^2 - b^2}}.$$

On the other hand we know that $\tilde{f}(t) \in (R_{a,b})$, and therefore

$$F(\tilde{f}) = \sqrt{a + \sqrt{a^2 - b^2}} \leq L(R_{a,b}).$$

PROOF OF (2.6). From $C_a \subset R_{a,b}$ and (2.9) we obtain that

$$L(C_a) \leq \sqrt{a + \sqrt{a^2 - b^2}}.$$

Letting $b \rightarrow a$ we obtain that

$$L(C_a) \leq \sqrt{a}.$$

However, in section 2 we found that for $\tilde{f}(t) = ae^{it}$ we have $F(\tilde{f}) = \sqrt{a}$.

REFERENCES

1. Landau, E. - *Einige Ungleichungen für zweimal differentiierbare Functionen*, Proc. London Math. Soc. (2), 13, 43-49 (1913).
2. Schoenberg, I. J. - *The elementary cases of Landau's problem of inequalities between derivatives*, Amer. Math. Monthly, 80, 121-158 (1973).
3. Schoenberg, I. J. - *The Landau problem for motions in a ring and in bounded continua*, Amer. Math. Monthly, 84, 1-12 (1977).